

Table S1

Means and Standard Deviations for Each Item From the Perceived Discrimination Measure (Wave 5)

	<i>M</i>	<i>SD</i>
Please determine how often you have experienced each event because of your RACE in the past 12 months...		
Being ignored, overlooked, or not given service (in a restaurant, store, etc.)	1.01	1.23
Being treated rudely or disrespectfully	1.13	1.24
Being accused of something or treated suspiciously	0.87	1.21
Others reacting to you as if they were afraid or intimidated	1.01	1.37
Being observed or followed while in public places	1.20	1.42
Being treated as if you were "stupid," being "talked down to"	0.97	1.31
Your ideas or opinions being minimized, ignored, or devalued	0.69	1.12
Overhearing or being told an offensive joke or comment	0.77	1.20
Being insulted, called a name, or harassed	0.37	1.09
Others expecting your work to be inferior	0.57	1.08
Not being taken seriously	1.00	1.34
Being left out of conversations or activities	0.69	1.17
Being treated in an "overly" friendly or superficial way	0.86	1.33
Other people avoiding you	0.53	1.12
Being mistaken for someone who serves others (i.e., janitor, bellboy, maid)	0.36	0.94
Being stared at by strangers	1.28	1.58
Being laughed at, made fun of, or taunted	0.36	0.98
Being mistaken for someone else of your same race	0.91	1.29
Being unfairly stopped, searched, questioned, physically threatened, or abused by the police	0.61	1.21
Not being hired for a job	0.90	1.27

Note. Scale ranged from 0 (*never happened to me*) to 5 (*once a week or more*).

Table S2

Correlations Among Exogenous Variables, Intercepts, Corresponding Intercept and Slope Factors, and Disturbances of Slope Factors

Variable	1	2	3	4	5	6	7	8	9	10	11	12
1. Male gender	—											
2. Highest grade completed	−.10*	—										
3. Family SES	.06	.21*	—									
4. Perceived discrimination intercept	.18*	.02	.02	—								
5. Perceived discrimination slope	—	—	—	−.01	—							
6. Anxiety symptoms intercept	−.10*	−.13*	−.12*	.29*	—	—						
7. Anxiety symptoms slope	—	—	—	—	.01	−.03	—					
8. Depressive symptoms intercept	−.13*	−.13*	−.11*	.29*	—	.72*	—	—				
9. Depressive symptoms slope	—	—	—	—	.14	—	.25*	−.03	—			
10. Cigarette use intercept	.10*	−.38*	−.06	.07	—	.15*	—	.15*	—	—		
11. Cigarette use slope	—	—	—	—	.12	—	.09*	—	.34*	−.10	—	
12. Alcohol use intercept	.16*	−.12*	.08	.23*	—	.08	—	.06	—	.42*	—	—
13. Alcohol use slope	—	—	—	—	.08	—	.10*	—	.35*	—	.45*	−.08

Note. SES = socioeconomic status.

* $p < .05$.