Table S1. Mean scores (s.d.) and results of ANOVAs for Beliefs about Medication by time and treatment group (n=58)

					Group x time	Group	Time
		Pre	Post	Follow-up	Pooled F(2,x)	Pooled F(1,x)	Pooled F(1,x)
BMQ Specific Necessity	Intervention	17.4 (3.0)	18.1 (2.6)	18.5 (2.5)	0.8, p=.460	0.1, p=.743	2.7, p=.066
	Control	18.2 (3.0)	17.9 (3.0)	18.5 (2.7)	0.0, p=. 100	0.1, p=.713	2.7, p=.000
BMQ Specific Concerns	Intervention	13.2 (3.2)	11.9 (3.7)	11.0 (2.5)	3.1, p=.047	.01, p=.924	9.2, p<.001
	Control	12.9 (3.4)	11.7 (3.7)	11.9 (3.0)	5.1, p=.017	.01, p=.921	7.2, p <.001
BMQ Necessity minus Concerns	Intervention	4.2 (4.5)	6.1 (4.9)	7.4 (3.4)	2.5, p=.081	0.02, p=.879	10.8, p<.001
	Control	5.3 (4.6)	6.2 (4.9)	6.6 (4.3)	2.3, p=.001	0.02, p=.077	10.0, p<.001

Note: df for F (i.e. x) are adjusted by inter-imputation variance to give pooled p-values and so vary by each statistic;

BMQ: Beliefs about Medication Questionnaire